

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## RH IL-17F (AF) SHD-100-152AF-100UG

Article Name	RH IL-17F (AF)
Biozol Catalog Number	SHD-100-152AF-100UG
Supplier Catalog Number	100-152AF-100UG
Alternative Catalog Number	SHD-100-152AF-100UG
Manufacturer	Fujifilm Irvine Scientific
Category	Proteine/Peptide
Application	Bioassay, SDS-PAGE
Species Reactivity	Human
Product Description	Interleukin 17F (IL-17F), a member of the IL-17 cytokine family, is secreted by activated CD4+ T cells and monocytes. IL-17F binds the IL-17 receptor related molecule, IL17RC, to promote the production of the interleukin 6 (IL-6), interleukin 8 (IL-8
Molecular Weight	15.0/30.1
UniProt	Q96PD4
Buffer	0.1% Trifluoroacetic Acid (TFA)
Source	E.coli
Purity	95
Form	Lyophilized from a 0.2 µm filtered solution

Sequence	MRKIPKVGHT FFQKPESCPP VPGGSMKLDI GIINENQRVS MSRNIESRST SPWNYTVTWD PNRYPSEVVQ AQCRNLGCIN AQGKEDISMN SVPIQQETLV VRRKHQGCSV SFQLEKVLVT VGCTCVTPVI HHVQ
Application Notes	Centrifuge vial before opening. Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.