



Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

RH MIG / CXCL9 SHD-100-184-100UG

| Article Name | RH MIG / CXCL9 |
|----------------------------|--|
| Biozol Catalog Number | SHD-100-184-100UG |
| Supplier Catalog Number | 100-184-100UG |
| Alternative Catalog Number | SHD-100-184-100UG |
| Manufacturer | Fujifilm Irvine Scientific |
| Category | Sonstiges |
| Application | Bioassay, SDS-PAGE |
| Species Reactivity | Human |
| Product Description | Monokine induced by gamma interferon (MIG or CXCL9) is a T cell chemoattractant during neuroinflammatory events. MIG production is stimulated by interferon gamma (IFN) and signals through the chemokine receptor CXCR3 |
| Molecular Weight | 11.7 |
| UniProt | Q07325 |
| Buffer | 0.1% Trifluoroacetic Acid (TFA) |
| Source | E.coli |
| Purity | 95 |
| Form | Lyophilized from a 0.2 μm filtered solution |

| Sequence | TPVVRKGRCS CISTNQGTIH LQSLKDLKQF APSPSCEKIE IIATLKNGVQ TCLNPDSADV KELIKKWEKQ VSQKKKQKNG KKHQKKKVLK VRKSQRSRQK KTT |
|-------------------|--|
| Application Notes | Centrifuge vial before opening. Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0. |