

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## RH NAP-2 SHD-100-210-100UG

Article Name	RH NAP-2
Biozol Catalog Number	SHD-100-210-100UG
Supplier Catalog Number	100-210-100UG
Alternative Catalog Number	SHD-100-210-100UG
Manufacturer	Fujifilm Irvine Scientific
Category	Sonstiges
Application	Bioassay, SDS-PAGE
Species Reactivity	Human
Product Description	Neutrophil activating peptide 2 (NAP-2), also known as CXCL7, is a member of the CXC family of chemokines. NAP-2 is a carboxylterminal fragment produced by proteolytic cleavage of the platelet basic protein (PBP). NAP-2 is released from platelets an
Molecular Weight	7.6
UniProt	P02775
Buffer	10 mM Sodium Phosphate, pH 7.5
Source	E.coli
Purity	95
Form	Lyophilized from a 0.2 µm filtered solution
Sequence	AELRCMCIKT TSGIHPKNIQ SLEVIGKGTH CNQVEVIATL KDGRKICLDP DAPRIKKIVQ KKLAGDESAD

## Application Notes

Centrifuge vial before opening. Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.