



Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

RM IL-17A (AF) SHD-200-59AF-100UG

Article Name	RM IL-17A (AF)
Biozol Catalog Number	SHD-200-59AF-100UG
Supplier Catalog Number	200-59AF-100UG
Alternative Catalog Number	SHD-200-59AF-100UG
Manufacturer	Fujifilm Irvine Scientific
Category	Proteine/Peptide
Application	Bioassay, SDS-PAGE
Species Reactivity	Mouse
Product Description	Interleukin 17A (IL-17A), also known as CTLA-8, is a member of the IL-17 family of proteins. IL-17A is a proinflammatory cytokine that is secreted by activated CD4+ and CD8+ T lymphocytes. IL-17A acts through its receptor, IL-17R, to promote increase
Molecular Weight	15.1/30.2
UniProt	Q62386
Buffer	0.1% Trifluoroacetic Acid (TFA)
Source	E.coli
Purity	95
Form	Lyophilized from a 0.2 μm filtered solution

Sequence	MAAIIPQSSA CPNTEAKDFL QNVKVNLKVF NSLGAKVSSR RPSDYLNRST SPWTLHRNED PDRYPSVIWE AQCRHQRCVN AEGKLDHHMN SVLIQQEILV LKREPESCPF TFRVEKMLVG VGCTCVASIV RQAA
Application Notes	Centrifuge vial before opening. Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.