

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

RM IL-17E (AF) SHD-200-60AF-100UG

Article Name	RM IL-17E (AF)
Biozol Catalog Number	SHD-200-60AF-100UG
Supplier Catalog Number	200-60AF-100UG
Alternative Catalog Number	SHD-200-60AF-100UG
Manufacturer	Fujifilm Irvine Scientific
Category	Proteine/Peptide
Application	Bioassay, SDS-PAGE
Species Reactivity	Mouse
Product Description	Interleukin 17E (IL-17E), also known as IL-25, is a member of the IL-17 family of cytokines. IL-17E binds to the IL-17RB receptor to stimulate the secretion of the proinflammatory interleukin 8 (IL-8) chemokine and to induce the activation of nuclear
Molecular Weight	17.6/35.2
UniProt	Q8VHC9
Buffer	0.1% Trifluoroacetic Acid (TFA)
Source	E.coli
Purity	90
Form	Lyophilized from a 0.2 µm filtered solution

Sequence	MVSLRIQEGC SHLPSCCPSK EQEPPEEWLK WSSASVSPPE PLSHTHHAES CRASKDGPLN SRAISPWSYE LDRDLNRVPQ DLYHARCLCP HCVSLQTGSH MDPLGNSVPL YHNQTVFYRR PCHGEEGTHR RYCLERRLYR VSLACVCVRP RVMA
Application Notes	Centrifuge vial before opening. Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.