

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Histone Antibody [1415-1], Unconjugated, Mouse, Monoclonal PRS-33-387

Article Name	Histone Antibody [1415-1], Unconjugated, Mouse, Monoclonal
Biozol Catalog Number	PRS-33-387
Supplier Catalog Number	33-387
Alternative Catalog Number	PRS-33-387-100
Manufacturer	ProSci
Host	Mouse
Category	Antikörper
Application	FC, IF, IHC-P, WB
Species Reactivity	Human, Mouse, Rat
Immunogen	Nuclei of human leukemia biopsy cells were used as the immunogen for the Histone antibody.
Conjugation	Unconjugated
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[1415-1]
NCBI	<a href="#">3005</a>
UniProt	<a href="#">P07305</a>
Buffer	PBS with 0.1 mg/ml BSA and 0.05% sodium azide
Form	Liquid

Application Dilute	Optimal dilutions for each application to be determined by the researcher
Application Notes	Western blot: 0.5-1 ug/ml Flow Cytometry: 0.5-1 ug/million cells in 0.1ml Immunofluorescence: 0.5-1 ug/ml Immunohistochemistry (FFPE): 0.5-1 ug/ml for 30 min at RT (1) Prediluted format : incubate for 30 min at RT (2) Optimal dilution of the Histone antibody should be determined by the researcher. 1. Staining of formalin/paraffin tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.