

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

IGF-BP3 Antibody, Unconjugated, Rabbit, Polyclonal PRS-38-250

| | |
|----------------------------|--|
| Article Name | IGF-BP3 Antibody, Unconjugated, Rabbit, Polyclonal |
| Biozol Catalog Number | PRS-38-250 |
| Supplier Catalog Number | 38-250 |
| Alternative Catalog Number | PRS-38-250-0.1 |
| Manufacturer | ProSci |
| Host | Rabbit |
| Category | Antikörper |
| Application | ELISA, WB |
| Species Reactivity | Human |
| Immunogen | Produced from sera of rabbits immunized with highly pure recombinant Rat SCF. Rat SCF specific antibody was purified by affinity chromatography employing an immobilized Rat SCF matrix. |
| Conjugation | Unconjugated |
| Clonality | Polyclonal |
| Concentration | batch dependent |
| NCBI | 3486 |
| UniProt | P17936 |
| Form | Lyophilized |
| Application Dilute | Centrifuge vial prior to opening. |

Application Notes

ELISA:Indirect:To detect hIGF-BP3 by indirect ELISA (using 100 μ L/well antibody solution) a concentration of 0.5 - 2.0 μ g/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hIGF-BP3.

SandwichTo detect hIGF-BP3 by sandwich ELISA (using 100 μ L/well antibody solution) a concentration of 0.5 - 2.0 μ g/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with our Biotinylated Anti-Human IGF-BP3 (38-153) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hIGF-BP3.

Western Blot:To detect hIGF-BP3 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 μ g/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIGF-BP3 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.