

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Flotillin 2 Peptide PRS-45-052P

Article Name	Flotillin 2 Peptide
Biozol Catalog Number	PRS-45-052P
Supplier Catalog Number	45-052P
Alternative Catalog Number	PRS-45-052P-0.1
Manufacturer	ProSci
Category	Proteine/Peptide
Application	Blocking
NCBI	2319
UniProt	Q14254
Buffer	Flotillin 2 peptide is supplied as 100 µg of dried peptide. When the peptide is reconstituted in 200 µL water, the concentration will be 0.5 mg/mL.
Form	Lyophilized

Application Notes

Flotillin 2 Peptide is used for blocking the activity of Flotillin 2 antibody (catalog number 45-052). The Flotillin 2 antibody ships at 0.5 mg/mL and the Flotillin 2 Peptide ships as a 100 ug pellet. When peptide is reconstituted in 200 uL water, the concentration will be 0.5 mg/mL, identical to that of the antibody. The best starting ratio would be 1:1 (which means molar excess of peptide relative to antibody when identical volumes are mixed). Mix equal volume of peptide and antibody at the required dilution and leave at ambient temperature for 20 minutes. Then incubate and develop in parallel two blots: one with the antibody and one with the antibody which has been pre-absorbed with the provided peptide.