



Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

FTCD Peptide PRS-45-057P

Article Name	FTCD Peptide
Biozol Catalog Number	PRS-45-057P
Supplier Catalog Number	45-057P
Alternative Catalog Number	PRS-45-057P-0.1
Manufacturer	ProSci
Category	Proteine/Peptide
Application	Blocking
NCBI	10841
UniProt	O95954
Buffer	FTCD peptide is supplied as 100 μg of dried peptide. When the peptide is reconstituted in 200 μL water, the concentration will be 0.5 mg/mL.
Form	Lyophilized

Application Notes

FTCD Peptide is used for blocking the activity of FTCD antibody (catalog number 45-057). The FTCD antibody ships at 0.5 mg/mL and the FTCD Peptide ships as a 100 ug pellet. When peptide is reconstituted in 200 uL water, the concentration will be 0.5 mg/mL, identical to that of the antibody. The best starting ratio would be 1:1 (which means molar excess of peptide relative to antibody when identical volumes are mixed). Mix equal volume of peptide and antibody at the required dilution and leave at ambient temperature for 20 minutes. Then incubate and develop in parallel two blots: one with the antibody and one with the antibody which has been preabsorbed with the provided peptide.