**BIOZOL**
Diagnostica Vertrieb GmbH, Oehleckerring 11-13
22419 Hamburg, Germany
**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99
**E-Mail:** info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

# Product Datasheet

## Clusterin Antibody [CLI-9], IgG1, Unconjugated, Mouse, Monoclonal PRS-XA-1007

| | |
|---|---|
| Article Name | Clusterin Antibody [CLI-9], IgG1, Unconjugated, Mouse, Monoclonal |
| Biozol Catalog Number | PRS-XA-1007 |
| Supplier Catalog Number | XA-1007 |
| Alternative Catalog Number | PRS-XA-1007-0.1 |
| Manufacturer | ProSci |
| Host | Mouse |
| Category | Antikörper |
| Application | ICC, IHC, IP, WB |
| Species Reactivity | Human |
| Immunogen | Clusterin (monoclonal CLI-9) antibody was raised against human clusterin (TRPM-2). |
| Conjugation | Unconjugated |
| Clonality | Monoclonal |
| Concentration | batch dependent |
| Clone Designation | [CLI-9] |
| Isotype | IgG1 |
| NCBI | 1191 |
| UniProt | P10909 |
| Buffer | PBS containing 0.02% sodium azide |

| | |
|---|---|
| Form | Liquid |
| Application Dilute | Optimal dilutions for each application to be determined by the researcher. |
| Application Notes | Clusterin antibody can be used for detection of human clusterin by immunocytochemistry, immunohistochemistry (frozen sections), immunoprecipitation, and western blot (1:2,000). Method: : Endogenous human Clusterin and recombinant FLAG-tagged Clusterin from cell lysates were resolved by SDS-PAGE under reducing conditions, transferred to nitrocellulose and incubated with a 1:2,000 dilution of the CL1-9 antibody on the left or with an anti-FLAG epitope-tag antibody on the right. Proteins were visualized using an HRP-conjugated anti-mouse IgG antibody and a chemiluminescence detection system. |