

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

APRIL Antibody [Aprily-1], IgG1, Unconjugated, Mouse, Monoclonal PRS-XA-1019

Article Name	APRIL Antibody [Aprily-1], IgG1, Unconjugated, Mouse, Monoclonal
Biozol Catalog Number	PRS-XA-1019
Supplier Catalog Number	XA-1019
Alternative Catalog Number	PRS-XA-1019-0.1
Manufacturer	ProSci
Host	Mouse
Category	Antikörper
Application	FC, WB
Species Reactivity	Human
Immunogen	APRIL (monoclonal Aprily-1) antibody was raised against recombinant human APRIL (aa. 93-233).
Conjugation	Unconjugated
Clonality	Monoclonal
Concentration	batch dependent
Clone Designation	[Aprily-1]
Isotype	IgG1
NCBI	8741
UniProt	O75888
Buffer	PBS containing 0.02% sodium azide

Form	Liquid
Application Dilute	Optimal dilutions for each application to be determined by the researcher.
Application Notes	<p>APRIL antibody can be used to recognize human APRIL by Flow Cytometry (also works for intracellular FC staining) and Western Blot. Anti-APRIL (MAb Aprily-1) does not cross react with mouse APRIL.</p> <p>Method: HEK 293 cells were mock transfected or transfected with an expression plasmid enabling surface expression of hAPRIL. Cells (5x10⁵) were incubated on ice for 30 min in 50 µl FACS buffer (PBS, 5% Fetal calf serum, 0.02% azide) containing 10 µg/mL of Aprily-1 antibody. After washing in FACS buffer, FITC-conjugated antibody to mouse IgG was added. Cells were incubated on ice for 30 min, washed and analyzed by flow cytometry.</p> <p>Method: Cell supernatants were first precleared with protein-A sepharose and then subjected to immunoprecipitation with rhBCMA:Fc and protein-A sepharose. Beads were washed twice with PBS. Immunoprecipitated materials were eluted with 2.5mM glycine pH 2.5 for 10 min, neutralized with Tris buffer and run on an SDS-PAGE gel and analyzed with 2 µg/mL Aprily-1 by Western blot. Note: In some migrations endogenous soluble APRIL resolved as a doublet around 20kDa (dendritic cells: DC). rhsMegaAPRIL served as a positive control.</p> <p>Method: 3 HEK 293T cells (5 x 10⁵) were mock transfected (thin line) or transfected with an expression plasmid enabling surface expression of mDR6 (thick line). Cells were incubated on ice for 30 min in 50 µl FACS buffer (PBS, 5% Fetal calf serum, 0.02% azide) containing 1 µg/mL of Luke-1 antibody. After washing in FACS buffer, PE-conjugated antibody to rat IgG was added. Cells were incubated on ice for 30 min, washed and then analyzed by flow cytometry.</p>