

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### IL-7 Antibody, Unconjugated, Rabbit, Polyclonal PRS-XP-5197

Article Name	IL-7 Antibody, Unconjugated, Rabbit, Polyclonal
Biozol Catalog Number	PRS-XP-5197
Supplier Catalog Number	XP-5197
Alternative Catalog Number	PRS-XP-5197-0.1
Manufacturer	ProSci
Host	Rabbit
Category	Antikörper
Application	ELISA, NeA, WB
Species Reactivity	Human
Immunogen	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hIL-7 (human Interleukin-7).
Conjugation	Unconjugated
Clonality	Polyclonal
Concentration	batch dependent
NCBI	<a href="#">3574</a>
UniProt	<a href="#">P13232</a>
Form	Lyophilized
Application Dilute	Centrifuge vial prior to opening.

Application Notes

Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of hIL-7 (1.00 ng/mL), a concentration of 0.016 - 0.03 µg/mL of this antibody is required. ELISA: To detect hIL-7 by direct ELISA (using 100 µL/well antibody solution) a concentration of at least 0.5 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4 ng/well of recombinant hIL-7. Sandwich: To detect hIL-7 by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.5 - 2.0 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with our Biotinylated Anti-Human IL-7 (XP-5197Bt) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hIL-7. Western Blot: To detect hIL-7 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIL-7 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.