

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## ECD Antibody / SGT1, Clone: [PCRP-ECD-1D10], Mouse, Monoclonal NSJ-V9605-20UG

ECD Antibody / SGT1, Clone: [PCRP-ECD-1D10], Mouse, Monoclonal
101 7/0001 30110
NSJ-V9605-20UG
/9605-20UG
NSJ-V9605-20UG
NSJ Bioreagents
Mouse
Antikörper
FACS, IF
Human
Recombinant full-length human protein was used as the immunogen for the ECD antibody.
nhibits MDM2-mediated degradation of p53/TP53 possibly by cooperating in part with TXNIP. May be involved transcriptional regulation. In vitro has intrinsic transactivation activity enhanced by EP300. May be a transcriptional activator required for
Monoclonal
PCRP-ECD-1D10]
095905
Protein A/G affinity

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Flow cytometry: 1-2ug/million cells,Immunofluorescence: 1-2ug/ml
Application Notes	Optimal dilution of the ECD antibody should be determined by the researcher.