

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Anti-Lck Monoclonal Antibody, Clone: [Clone: ADBE-12], Rabbit BOB-M00425

|                            |  |
|----------------------------|--|
| Article Name               | Anti-Lck Monoclonal Antibody, Clone: [Clone: ADBE-12], Rabbit  |
| Biozol Catalog Number      | BOB-M00425   |
| Supplier Catalog Number    | M00425   |
| Alternative Catalog Number | BOB-M00425-100UL,BOB-M00425-30UL   |
| Manufacturer               | Boster Bio   |
| Host                       | Rabbit   |
| Category                   | Antikörper   |
| Application                | FC, ICC, IF, IHC, IP, WB   |
| Species Reactivity         | Human  |
| Immunogen                  | A synthesized peptide derived from human Lck The Src family of protein tyrosine kinases, which includes Src, Lyn, Fyn, Yes, Lck, Blk, and Hck, are important in the regulation of growth and differentiation of eukaryotic cells. Src activity is regulated by tyrosine phosphorylation at two sites, but with opposing effects. |
| Product Description        | Boster Bio Anti-Lck Monoclonal Antibody catalog M00425. Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human....  |
| Clonality                  | Monoclonal   |
| Concentration              | 0.5mg/ml   |
| Clone Designation          | [Clone: ADBE-12]   |
| Molecular Weight           | 11 kDa   |

|                    |   |
|--------------------|---|
| Sensitivity        | <20pg/ml  |
| UniProt            | <a href="#">P06239</a>  |
| Buffer             | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA. |
| Purity             | Affinity-chromatography   |
| Form               | Liquid  |
| Application Dilute | WB 1:500-2000IHC 1:50-200ICC/IF 1:50-200IP 1:20FC 1:20  |