

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Human IgM Immunoglobulin(IM373), CF405S conjugate, 0.1mg/mL, Clone: [IM373], Mouse, Monoclonal BOT-BNC040373-100

| | |
|----------------------------|---|
| Article Name | Human IgM Immunoglobulin(IM373), CF405S conjugate, 0.1mg/mL, Clone: [IM373], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC040373-100 |
| Supplier Catalog Number | BNC040373-100 |
| Alternative Catalog Number | BOT-BNC040373-100-100UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length heavy chain of human IgM |
| Conjugation | CF405S |
| Product Description | Recognizes a protein of 75 kDa, identified as mu heavy chain of human immunoglobulins. It does not cross-react with alpha (IgA), gamma (IgG), epsilon (IgE), or delta (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes. This MAb is ... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [IM373] |

| | |
|-------------------|---|
| Molecular Weight | 50-75 kDa |
| UniProt | P01871 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |