

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Neurofilament (NF-H)(NF421), CF405S conjugate, 0.1mg/mL, IgG1, Clone: [NF421], Mouse, Monoclonal BOT-BNC040421-100

| | |
|----------------------------|---|
| Article Name | Neurofilament (NF-H)(NF421), CF405S conjugate, 0.1mg/mL, IgG1, Clone: [NF421], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC040421-100 |
| Supplier Catalog Number | BNC040421-100 |
| Alternative Catalog Number | BOT-BNC040421-100-100UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, IF, IHC, WB |
| Species Reactivity | Gallus, Human, Mouse, Porcine, Rat |
| Immunogen | Human recombinant neurofilament protein |
| Conjugation | CF405S |
| Product Description | This MAb reacts with a 200 kDa protein, identified as heavy sub-unit of neurofilaments (NF-H). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neu... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [NF421] |

| | |
|-------------------|--|
| Molecular Weight | 200 kDa |
| Isotype | IgG1 |
| UniProt | P12036 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |