

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

FGF23(FGF23/638), CF405S conjugate, 0.1mg/mL, Clone: [FGF23/638], Mouse, Monoclonal BOT-BNC040638-500

| | |
|----------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name | FGF23(FGF23/638), CF405S conjugate, 0.1mg/mL, Clone: [FGF23/638], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC040638-500 |
| Supplier Catalog Number | BNC040638-500 |
| Alternative Catalog Number | BOT-BNC040638-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Species Reactivity | Human |
| Immunogen | Recombinant human FGF23 protein |
| Conjugation | CF405S |
| Product Description | Fibroblast growth factor-1 (FGF-1, acidic FGF) and fibroblast growth factor-2 (FGF-2, basic FGF) are members of a family of growth factors that stimulate proliferation of cells of mesenchymal, epithelial and neuroectodermal origin. Additional members ... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [FGF23/638] |
| Molecular Weight | 12-32 kDa |

| | |
|-------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| UniProt | Q9GZV9 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |