

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

bcl-2(BCL2/796), CF405S conjugate, 0.1mg/mL, Clone: [BCL2/796], Mouse, Monoclonal BOT-BNC040796-100

Article Name	bcl-2(BCL2/796), CF405S conjugate, 0.1mg/mL, Clone: [BCL2/796], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC040796-100
Supplier Catalog Number	BNC040796-100
Alternative Catalog Number	BOT-BNC040796-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC, WB
Species Reactivity	Human
Immunogen	Recombinant full-length human bcl-2 protein
Conjugation	CF405S
Product Description	This antibody recognizes a protein of 25-26 kDa, identified as Bcl-2. This antibody shows no cross-reaction with Bcl-x or Bax proteins. Bcl-2 suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural ...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[BCL2/796]

Molecular Weight	25-26 kDa
UniProt	P10415
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user