

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Cytokeratin 19(KRT19/800), CF405S conjugate, 0.1mg/mL, Clone: [KRT19/800], Mouse, Monoclonal BOT-BNC040800-100**

|                            |   |
|----------------------------|---|
| Article Name               | Cytokeratin 19(KRT19/800), CF405S conjugate, 0.1mg/mL, Clone: [KRT19/800], Mouse, Monoclonal  |
| Biozol Catalog Number      | BOT-BNC040800-100   |
| Supplier Catalog Number    | BNC040800-100   |
| Alternative Catalog Number | BOT-BNC040800-100-100UL   |
| Manufacturer               | Biotium   |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | IF, WB  |
| Species Reactivity         | Human, Mouse, Rat   |
| Immunogen                  | Recombinant human KRT19 protein   |
| Conjugation                | CF405S  |
| Product Description        | Recognizes a protein of 40 kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epit... |
| Clonality                  | Monoclonal  |
| Concentration              | 0.1 mg/mL   |
| Clone Designation          | [KRT19/800]   |

|                   |  |
|-------------------|--|
| Molecular Weight  | 40 kDa   |
| UniProt           | <a href="#">P08727</a>   |
| Buffer            | PBS, 0.1% BSA, 0.05% azide   |
| Source            | Animal   |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |