

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Ep-CAM / CD326(EGP40/837), CF405S conjugate, 0.1mg/mL, Clone: [EGP40/837], Mouse, Monoclonal BOT-BNC040837-100

| | |
|----------------------------|---|
| Article Name | Ep-CAM / CD326(EGP40/837), CF405S conjugate, 0.1mg/mL, Clone: [EGP40/837], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC040837-100 |
| Supplier Catalog Number | BNC040837-100 |
| Alternative Catalog Number | BOT-BNC040837-100-100UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, IF, IHC, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant human TACSTD1 protein |
| Conjugation | CF405S |
| Product Description | EGP40 is a 40-43 kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majo... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [EGP40/837] |

| | |
|-------------------|--|
| Molecular Weight | 40-43 kDa |
| UniProt | P16422 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Does not react with mouse or rat, others not known Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |