

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Glypican-3(1G12), CF405S conjugate, 0.1mg/mL, Clone: [1G12], Mouse, Monoclonal BOT-BNC040862-100**

|                            |   |
|----------------------------|---|
| Article Name               | Glypican-3(1G12), CF405S conjugate, 0.1mg/mL, Clone: [1G12], Mouse, Monoclonal  |
| Biozol Catalog Number      | BOT-BNC040862-100   |
| Supplier Catalog Number    | BNC040862-100   |
| Alternative Catalog Number | BOT-BNC040862-100-100UL   |
| Manufacturer               | Biotium   |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | FC, IF, IHC, WB   |
| Species Reactivity         | Human   |
| Immunogen                  | A recombinant fragment containing amino acids 511-580 of human glypican-3   |
| Conjugation                | CF405S  |
| Product Description        | Glypican-3 (GPC3) is a glycosylphosphatidylinositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, mel... |
| Clonality                  | Monoclonal  |
| Concentration              | 0.1 mg/mL   |
| Clone Designation          | [1G12]  |

|                   |   |
|-------------------|---|
| Molecular Weight  | 67 kDa  |
| UniProt           | <a href="#">P51654</a>  |
| Buffer            | PBS, 0.1% BSA, 0.05% azide  |
| Source            | Animal  |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |