

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **MHC II (HLA-DRA)(IPO-10), CF405S conjugate, 0.1mg/mL, Clone: [IPO-10], Mouse, Monoclonal BOT-BNC041090-500**

Article Name	MHC II (HLA-DRA)(IPO-10), CF405S conjugate, 0.1mg/mL, Clone: [IPO-10], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC041090-500
Supplier Catalog Number	BNC041090-500
Alternative Catalog Number	BOT-BNC041090-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Species Reactivity	Human, Monkey
Immunogen	Spleen cells of a patient with hairy cell leukemia (Daudi cells)
Conjugation	CF405S
Product Description	MAb IPO-10 defines the antigen that appears on B cell progenitors following HLA-DR and preceding CD10, CD19, CD22, CD37 and cym. It is expressed on resting B cells and then reappears and persists in cytoplasm and on cell surface until cytoplasmic Ig ...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[IPO-10]
Molecular Weight	36 kDa

UniProt	<a href="#">P01903</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry