

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **PTH / Parathyroid Hormone (N-Terminal)(PTH/1173), CF405S conjugate, 0.1mg/mL, Clone: [PTH/1173], Mouse, Monoclonal BOT-BNC041173-100**

|                            |   |
|----------------------------|---|
| Article Name               | PTH / Parathyroid Hormone (N-Terminal)(PTH/1173), CF405S conjugate, 0.1mg/mL, Clone: [PTH/1173], Mouse, Monoclonal  |
| Biozol Catalog Number      | BOT-BNC041173-100   |
| Supplier Catalog Number    | BNC041173-100   |
| Alternative Catalog Number | BOT-BNC041173-100-100UL   |
| Manufacturer               | Biotium   |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | IHC   |
| Species Reactivity         | Human   |
| Immunogen                  | Recombinant fragment (84 amino acid residues from C-terminus) of human PTH protein  |
| Conjugation                | CF405S  |
| Product Description        | The epitope of this MAb maps in the C-terminus of PTH, a hormone produced by the parathyroid gland that regulates the concentration of calcium and phosphorus in extracellular fluid. This hormone elevates blood calcium levels by dissolving the salts i... |
| Clonality                  | Monoclonal  |
| Concentration              | 0.1 mg/mL   |
| Clone Designation          | [PTH/1173]  |

|                   |  |
|-------------------|--|
| Molecular Weight  | 9 kDa  |
| UniProt           | <a href="#">P01270</a>   |
| Buffer            | PBS, 0.1% BSA, 0.05% azide   |
| Source            | Animal   |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |