

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Cytokeratin 18 (KRT18)(B23.1), CF405S conjugate, 0.1mg/mL, Clone: [B23.1], Mouse, Monoclonal BOT-BNC041263-100**

Article Name	Cytokeratin 18 (KRT18)(B23.1), CF405S conjugate, 0.1mg/mL, Clone: [B23.1], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC041263-100
Supplier Catalog Number	BNC041263-100
Alternative Catalog Number	BOT-BNC041263-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC, WB
Species Reactivity	Mouse
Immunogen	PMC-42 human breast carcinoma cells
Conjugation	CF405S
Product Description	This MAAb reacts with a wide variety of simple epithelia. It does not react with stratified squamous epithelia. It reacts with epithelial tumors of the gastrointestinal tract, lung, breast, pancreas, ovary, and thyroid. Cytokeratin 18, which belongs to...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[B23.1]

Molecular Weight	45 kDa
UniProt	<a href="#">P05783</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	<p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 min at RT Flow cytometry: 0.5-1 ug/million cells Western Blot 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user</p>