

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

c-Myc Oncoprotein(MYC275 + MYC909), CF405S conjugate, 0.1mg/mL, Clone: [MYC275 MYC909], Mouse, Monoclonal BOT-BNC041269-100

| | |
|----------------------------|---|
| Article Name | c-Myc Oncoprotein(MYC275 + MYC909), CF405S conjugate, 0.1mg/mL, Clone: [MYC275 MYC909], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC041269-100 |
| Supplier Catalog Number | BNC041269-100 |
| Alternative Catalog Number | BOT-BNC041269-100-100UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant human c-myc protein |
| Conjugation | CF405S |
| Product Description | This antibody recognizes a transcription factor of 64-67 kDa, identified as c-myc. This MAb shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or over-expressed in a var... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [MYC275 MYC909] |

| | |
|-------------------|--|
| Molecular Weight | 62-64 kDa |
| UniProt | P01106 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at RT Flow cytometry: 0.5-1 ug/million cells Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |