

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

HPV-16 E6 + HPV-18 E6 (Human Papilloma Virus 16/18)(HPV16/1295 + HPV18/1297), CF405S conjugate, 0.1mg/mL, Clone: [HPV16/1295 HPV18/1297], Mouse, Monoclonal BOT-BNC041340-500

| | |
|----------------------------|---|
| Article Name | HPV-16 E6 + HPV-18 E6 (Human Papilloma Virus 16/18)(HPV16/1295 + HPV18/1297), CF405S conjugate, 0.1mg/mL, Clone: [HPV16/1295 HPV18/1297], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC041340-500 |
| Supplier Catalog Number | BNC041340-500 |
| Alternative Catalog Number | BOT-BNC041340-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Species Reactivity | Virus |
| Immunogen | HPV18 E6-beta-galactosidase fusion protein |
| Conjugation | CF405S |
| Product Description | Human papilloma viruses (HPVs) can be classified as either high risk or low risk according to their association with cancer. HPV16 and HPV18 are the most common of the high risk group while HPV6 and HPV11 are among the low risk types. Approximately 9... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [HPV16/1295 HPV18/1297] |

| | |
|-------------------|--|
| Molecular Weight | 16/17 kDa |
| UniProt | Not Applicable |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |