

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD257 / BAFF / TNFSF13B (C257/1638), CF405S conjugate, 0.1mg/mL, IgG1, Clone: [C257/1638], Mouse, Monoclonal BOT-BNC041638-500

| | |
|----------------------------|---|
| Article Name | CD257 / BAFF / TNFSF13B (C257/1638), CF405S conjugate, 0.1mg/mL, IgG1, Clone: [C257/1638], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC041638-500 |
| Supplier Catalog Number | BNC041638-500 |
| Alternative Catalog Number | BOT-BNC041638-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Species Reactivity | Human |
| Immunogen | Purified recombinant human full-length CD257 protein |
| Conjugation | CF405S |
| Product Description | BAFF/CD257 is a type II transmembrane protein, and a member of the tumor necrosis factor ligand superfamily. It is proteolytically cleaved to form a soluble protein. BAFF, like other TNF ligand family members, forms homo-trimers. The predicted molecu... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [C257/1638] |
| Molecular Weight | 31 kDa (Monomer) |

| | |
|-------------------|--|
| Isotype | IgG1 |
| UniProt | Q9Y275 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |