

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD86 (Dendritic Cells Maturation Marker) (rC86/1146), CF405S conjugate, 0.1mg/mL, Clone: [rC86/1146], Mouse, Monoclonal BOT-BNC042083-500

| | |
|----------------------------|---|
| Article Name | CD86 (Dendritic Cells Maturation Marker) (rC86/1146), CF405S conjugate, 0.1mg/mL, Clone: [rC86/1146], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC042083-500 |
| Supplier Catalog Number | BNC042083-500 |
| Alternative Catalog Number | BOT-BNC042083-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, IF, IHC |
| Species Reactivity | Human, Mouse, Rat |
| Immunogen | Recombinant full-length human CD86 protein |
| Conjugation | CF405S |
| Product Description | This antibody recognizes a protein of 70 kDa, which is identified as CD86. CD86 is a type I transmembrane glycoprotein and a member of the immunoglobulin superfamily of cell surface receptors. It is expressed at high levels on resting peripheral mono... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [rC86/1146] |

| | |
|-------------------|---|
| Molecular Weight | 70 kDa |
| UniProt | P42081 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 2-4 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris buffer with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |