

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Cytokeratin 8 (KRT8) (KRT8/2174R), CF405S conjugate, 0.1mg/mL, Clone: [KRT8/2174R], Rabbit, Monoclonal BOT-BNC042174-100**

|                            |                                                                                                                                                                                                                                                               |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name               | Cytokeratin 8 (KRT8) (KRT8/2174R), CF405S conjugate, 0.1mg/mL, Clone: [KRT8/2174R], Rabbit, Monoclonal                                                                                                                                                        |
| Biozol Catalog Number      | BOT-BNC042174-100                                                                                                                                                                                                                                             |
| Supplier Catalog Number    | BNC042174-100                                                                                                                                                                                                                                                 |
| Alternative Catalog Number | BOT-BNC042174-100-100UL                                                                                                                                                                                                                                       |
| Manufacturer               | Biotium                                                                                                                                                                                                                                                       |
| Host                       | Rabbit                                                                                                                                                                                                                                                        |
| Category                   | Antikörper                                                                                                                                                                                                                                                    |
| Application                | IHC, WB                                                                                                                                                                                                                                                       |
| Species Reactivity         | Bovine, Human, Porcine, Rabbit, Sheep                                                                                                                                                                                                                         |
| Immunogen                  | Recombinant full-length human KRT8 protein                                                                                                                                                                                                                    |
| Conjugation                | CF405S                                                                                                                                                                                                                                                        |
| Product Description        | Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). CK8 is primarily found in the non-squamous epithelia and is present in majority of adenoc... |
| Clonality                  | Monoclonal                                                                                                                                                                                                                                                    |
| Concentration              | 0.1 mg/mL                                                                                                                                                                                                                                                     |
| Clone Designation          | [KRT8/2174R]                                                                                                                                                                                                                                                  |

|                   |                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|-------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molecular Weight  | 52.5 kDa                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| UniProt           | <a href="#">P05787</a>                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Buffer            | PBS, 0.1% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Source            | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1.0 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |