

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Bcl-2 (Apoptosis & Follicular Lymphoma Marker) (BCL2/2210R), CF405S conjugate, 0.1mg/mL, Clone: [BCL2/2210R], Rabbit, Monoclonal BOT-BNC042210-100**

Article Name	Bcl-2 (Apoptosis & Follicular Lymphoma Marker) (BCL2/2210R), CF405S conjugate, 0.1mg/mL, Clone: [BCL2/2210R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNC042210-100
Supplier Catalog Number	BNC042210-100
Alternative Catalog Number	BOT-BNC042210-100-100UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	IHC, WB
Species Reactivity	Human
Immunogen	Recombinant full-length human bcl-2 protein
Conjugation	CF405S
Product Description	This antibody recognizes a protein of 25-26 kDa, identified as Bcl-2. This antibody shows no cross-reaction with Bcl-x or Bax proteins. Bcl-2 suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural ...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[BCL2/2210R]

Molecular Weight	25-26 kDa
UniProt	<a href="#">P10415</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris with 1 mM EDTA Buffer pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user