

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **ACTH (Adrenocorticotrophic Hormone) (Pituitary Marker) (rCLIP/1407), CF405S conjugate, 0.1mg/mL, IgG1, Clone: [rCLIP/1407], Mouse, Monoclonal BOT-BNC042296-500**

Article Name	ACTH (Adrenocorticotrophic Hormone) (Pituitary Marker) (rCLIP/1407), CF405S conjugate, 0.1mg/mL, IgG1, Clone: [rCLIP/1407], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC042296-500
Supplier Catalog Number	BNC042296-500
Alternative Catalog Number	BOT-BNC042296-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human, Mouse, Rat
Immunogen	Synthetic peptide corresponding to (aa 162-176) of human ACTH (NGAEDESAEAFPLEF)
Conjugation	CF405S
Product Description	ACTH (same as Corticotropin) is a 39 amino acid active peptide produced by the anterior pituitary. This MAb is specific to CLIP (aa25-39 of ACTH), does not react with Synacthen (aa1-24 of ACTH). POMC (pro-opiomelanocortin or corticotropin-lipotropin)...
Clonality	Monoclonal
Concentration	0.1 mg/mL

Clone Designation	[rCLIP/1407]
Molecular Weight	ACTH is ~5 kDa, and the POMC precursor is ~30 kDa. The molecular weight of POMC depends upon isoform variation and post-translational modifications.
Isotype	IgG1
UniProt	<a href="#">P01189</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user