

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **CD269 / TNFRSF17 / BCMA (B-Cell Maturation Protein) (BCMA/2366), CF405S conjugate, 0.1mg/mL, Clone: [BCMA/2366], Mouse, Monoclonal BOT-BNC042366-100**

Article Name	CD269 / TNFRSF17 / BCMA (B-Cell Maturation Protein) (BCMA/2366), CF405S conjugate, 0.1mg/mL, Clone: [BCMA/2366], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC042366-100
Supplier Catalog Number	BNC042366-100
Alternative Catalog Number	BOT-BNC042366-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant human CD269 protein fragment (around aa 78-184) (exact sequence is proprietary)
Conjugation	CF405S
Product Description	The B cell maturation protein (BCMA) is a type I integral membrane protein that belongs to the tumor necrosis factor receptor (TNF-R) superfamily. It is expressed as a 184 amino acid peptide that is expressed only in mature B-lymphocytes and is local...
Clonality	Monoclonal
Concentration	0.1 mg/mL

Clone Designation	[BCMA/2366]
Molecular Weight	20 kDa
UniProt	<a href="#">Q02223</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: For coating order antibody without BSA Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user