

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cytokeratin 6A (KRT6A) (Basal Cell Marker) (KRT6A/2368), CF405S conjugate, 0.1mg/mL, Clone: [KRT6A/2368], Mouse, Monoclonal BOT-BNC042368-100

| | |
|----------------------------|---|
| Article Name | Cytokeratin 6A (KRT6A) (Basal Cell Marker) (KRT6A/2368), CF405S conjugate, 0.1mg/mL, Clone: [KRT6A/2368], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC042368-100 |
| Supplier Catalog Number | BNC042368-100 |
| Alternative Catalog Number | BOT-BNC042368-100-100UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length human Cytokeratin 6A (KRT6A) protein |
| Conjugation | CF405S |
| Product Description | This MAb recognizes a protein of 56 kDa, identified as cytokeratin 6A (KRT6A). In humans, multiple isoforms of Cytokeratin 6 (6A-6F), encoded by several highly homologous genes, have distinct tissue expression patterns. Cytokeratin 6A is the dominant... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [KRT6A/2368] |

| | |
|-------------------|---|
| Molecular Weight | 56 kDa |
| UniProt | P02538 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |