

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD3e (T-Cell Marker) (C3e/2478), CF405S conjugate, 0.1mg/mL, Clone: [C3e/2478], Mouse, Monoclonal BOT-BNC042478-500

| | |
|----------------------------|---|
| Article Name | CD3e (T-Cell Marker) (C3e/2478), CF405S conjugate, 0.1mg/mL, Clone: [C3e/2478], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC042478-500 |
| Supplier Catalog Number | BNC042478-500 |
| Alternative Catalog Number | BOT-BNC042478-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, WB |
| Species Reactivity | Human, Primate |
| Immunogen | Recombinant human CD3e extracellular domain fragment (around aa 23-119) (exact sequence is proprietary) |
| Conjugation | CF405S |
| Product Description | The immunogen for this Mab is derived from the extracellular domain of CD3 epsilon chain. CD3 consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28 kDa. The CD3 complex is closel... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [C3e/2478] |

| | |
|-------------------|--|
| Molecular Weight | 20 kDa |
| UniProt | P07766 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: 2-4 ug/m for coating order Ab without BSA Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |