

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

PD-L1 / PDCD1LG1 / CD274 / B7-H1 (Cancer Immunotherapy Target)(PDL1/2741), CF405S conjugate, 0.1mg/mL, Clone: [PDL1/2741], Mouse, Monoclonal BOT-BNC042741-500

| | |
|----------------------------|---|
| Article Name | PD-L1 / PDCD1LG1 / CD274 / B7-H1 (Cancer Immunotherapy Target)(PDL1/2741), CF405S conjugate, 0.1mg/mL, Clone: [PDL1/2741], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC042741-500 |
| Supplier Catalog Number | BNC042741-500 |
| Alternative Catalog Number | BOT-BNC042741-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Species Reactivity | Human |
| Immunogen | Recombinant fragment of human CD274 protein (around aa39-191) (exact sequence is proprietary) |
| Conjugation | CF405S |
| Product Description | PD-L1 is a checkpoint regulator in immune cells, it is expressed on immune or non-hematopoietic cells. Expression of the protein is seen during pregnancy where it has a role in suppressing the immune system. PD-L1 induces an inhibitory signal in acti... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [PDL1/2741] |

| | |
|-------------------|--|
| Molecular Weight | 37-50 kDa |
| UniProt | Q9NZQ7 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |