

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

ATRX / RAD54 (Alpha Thalassemia Mental Retardation) (ATRX/2900R), CF405S conjugate, 0.1mg/mL, Clone: [ATRX/2900R], Rabbit, Monoclonal BOT-BNC042900-100

| | |
|----------------------------|---|
| Article Name | ATRX / RAD54 (Alpha Thalassemia Mental Retardation) (ATRX/2900R), CF405S conjugate, 0.1mg/mL, Clone: [ATRX/2900R], Rabbit, Monoclonal |
| Biozol Catalog Number | BOT-BNC042900-100 |
| Supplier Catalog Number | BNC042900-100 |
| Alternative Catalog Number | BOT-BNC042900-100-100UL |
| Manufacturer | Biotium |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length human ATRX protein |
| Conjugation | CF405S |
| Product Description | ATRX is a member of the Snf2 family of helicase/ATPases, which contribute to the remodeling of the nucleosome structure in an ATP-dependent manner, and facilitate the initiation of transcription and replication. Structurally, ATRX contains a PHD zinc... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [ATRX/2900R] |

| | |
|-------------------|---|
| Molecular Weight | 280 kDa |
| UniProt | P46100 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris with 1 mM EDTA Buffer pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |