

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **GFAP(GA-5), CF640R conjugate, 0.1mg/mL, IgG1, Clone: [GA-5], Mouse, Monoclonal BOT-BNC400167-500**

Article Name	GFAP(GA-5), CF640R conjugate, 0.1mg/mL, IgG1, Clone: [GA-5], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC400167-500
Supplier Catalog Number	BNC400167-500
Alternative Catalog Number	BOT-BNC400167-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	FC, IF, IHC, WB
Species Reactivity	Bovine, Gallus, Human, Mouse, Porcine, Rabbit, Rat
Immunogen	GFAP isolated from pig spinal cord
Conjugation	CF640R
Product Description	This MAAb recognizes a protein of ~50 kDa which is identified as Glial Fibrillary Acidic Protein (GFAP). It shows no cross-reaction with other intermediate filament proteins. GFAP is specifically found in astroglia. GFAP is a very popular marker for I...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[GA-5]

Molecular Weight	~50 kDa
Isotype	IgG1
UniProt	<a href="#">P14136</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	<p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user</p>