

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **CD45RB(PD7/26), CF640R conjugate, 0.1mg/mL, Clone: [PD7/26], Mouse, Monoclonal BOT-BNC400472-500**

|                            |   |
|----------------------------|---|
| Article Name               | CD45RB(PD7/26), CF640R conjugate, 0.1mg/mL, Clone: [PD7/26], Mouse, Monoclonal  |
| Biozol Catalog Number      | BOT-BNC400472-500   |
| Supplier Catalog Number    | BNC400472-500   |
| Alternative Catalog Number | BOT-BNC400472-500-500UL   |
| Manufacturer               | Biotium   |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | FC, IHC, WB   |
| Species Reactivity         | Human   |
| Immunogen                  | Human peripheral blood lymphocytes maintained in T cell growth factor   |
| Conjugation                | CF640R  |
| Product Description        | CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms ar... |
| Clonality                  | Monoclonal  |
| Concentration              | 0.1 mg/mL   |

|                   |   |
|-------------------|---|
| Clone Designation | [PD7/26]  |
| Molecular Weight  | 180-220 kDa   |
| UniProt           | <a href="#">P08575</a>  |
| Buffer            | PBS, 0.1% BSA, 0.05% azide  |
| Source            | Animal  |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |