

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD45 / LCA(2B11 + PD7/26), CF640R conjugate, 0.1mg/mL, Clone: [2B11 PD7/26], Mouse, Monoclonal BOT-BNC400745-500

| | |
|----------------------------|---|
| Article Name | CD45 / LCA(2B11 + PD7/26), CF640R conjugate, 0.1mg/mL, Clone: [2B11 PD7/26], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC400745-500 |
| Supplier Catalog Number | BNC400745-500 |
| Alternative Catalog Number | BOT-BNC400745-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, IF, IHC, IP |
| Species Reactivity | Canine, Human |
| Immunogen | Isolated neoplastic cells from T cell lymphoma (2B11), and human peripheral blood lymphocytes maintained in T cell growth factor (PD7/26) |
| Conjugation | CF640R |
| Product Description | CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms ar... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |

| | |
|-------------------|--|
| Clone Designation | [2B11 PD7/26] |
| Molecular Weight | 180-220 kDa |
| UniProt | P08575 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |