

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Phosphotyrosine(PY793), CF640R conjugate, 0.1mg/mL, IgG2b, Clone: [PY793], Mouse, Monoclonal BOT-BNC400793-500

| | |
|----------------------------|---|
| Article Name | Phosphotyrosine(PY793), CF640R conjugate, 0.1mg/mL, IgG2b, Clone: [PY793], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC400793-500 |
| Supplier Catalog Number | BNC400793-500 |
| Alternative Catalog Number | BOT-BNC400793-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | All |
| Immunogen | Phosphotyrosine conjugated to BSA |
| Conjugation | CF640R |
| Product Description | Protein phosphorylation is a fundamental event in the regulation of a large number of intracellular processes. Phosphorylation of specific tyrosine residues is the result of activation or stimulation of their respective protein tyrosine kinases. The ... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [PY793] |

| | |
|-------------------|--|
| Molecular Weight | Depends upon the phosphorylated target |
| Isotype | IgG2b |
| UniProt | Not Applicable |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 1-2 ug/mL No pretreatment is required for staining of formalin-fixed, paraffin-embedded tissues Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |