

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Cytokeratin, Acidic (Type I or LMW) (Epithelial Marker) (KRTL/1377), CF640R conjugate, 0.1mg/mL, IgG1, Clone: [KRTL/1377], Mouse, Monoclonal BOT-BNC401377-100**

Article Name	Cytokeratin, Acidic (Type I or LMW) (Epithelial Marker) (KRTL/1377), CF640R conjugate, 0.1mg/mL, IgG1, Clone: [KRTL/1377], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC401377-100
Supplier Catalog Number	BNC401377-100
Alternative Catalog Number	BOT-BNC401377-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Species Reactivity	Human
Immunogen	Recombinant human KRT77 protein fragment (exact sequence is proprietary)
Conjugation	CF640R
Product Description	This MAbs recognizes the 56.5 kDa (CK10), 50 kDa (CK14), 50 kDa (CK15), 48 kDa (CK16), 40 kDa (CK19) keratins of the acidic (Type I or LMW) subfamily. Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) sub...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[KRTL/1377]

Molecular Weight	56.5 kDa (CK10), 50 kDa (CK14), 50 kDa (CK15), 48 kDa (CK16), 40 kDa (CK19)
Isotype	IgG1
UniProt	<a href="#">Q7Z794</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry