

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **TIMP2 (Tissue Inhibitor of Metalloproteinase 2) (TIMP2/2044), CF640R conjugate, 0.1mg/mL, Clone: [TIMP2/2044], Mouse, Monoclonal BOT-BNC402044-100**

Article Name	TIMP2 (Tissue Inhibitor of Metalloproteinase 2) (TIMP2/2044), CF640R conjugate, 0.1mg/mL, Clone: [TIMP2/2044], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC402044-100
Supplier Catalog Number	BNC402044-100
Alternative Catalog Number	BOT-BNC402044-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant human TIMP2 protein fragment (aa 48-220) (exact sequence is proprietary)
Conjugation	CF640R
Product Description	This antibody recognizes a protein of 21 kDa, identified as tissue inhibitor of metalloproteinases-2 (TIMP-2). It is closely related to TIMP-1 and shows the highest binding affinity to both the latent (pro) and active forms of 72 kDa Type IV collagen...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[TIMP2/2044]

Molecular Weight	21 kDa
UniProt	<a href="#">P16035</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: 2-4 ug/m for coating order Ab without BSA Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user