

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **bcl-x(BX006), CF647 conjugate, 0.1mg/mL, IgG2a, Clone: [BX006], Mouse, Monoclonal BOT-BNC470006-100**

Article Name	bcl-x(BX006), CF647 conjugate, 0.1mg/mL, IgG2a, Clone: [BX006], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC470006-100
Supplier Catalog Number	BNC470006-100
Alternative Catalog Number	BOT-BNC470006-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC, WB
Species Reactivity	Human, Mouse, Porcine, Rat
Immunogen	A synthetic peptide, aa 3-14 (Cys-QSNRELVDFLS) of human Bcl-X protein.
Conjugation	CF647
Product Description	Bcl-x belongs to the Bcl-2 family of proteins and plays a crucial role in apoptosis. Bcl-X has two isoforms: Bcl-XL (long), a 26 kDa protein which suppresses cell death, and Bcl-XS (short), a 178 amino acid protein lacking a 63 amino acid domain whic...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[BX006]

Molecular Weight	27 kDa
Isotype	IgG2a
UniProt	<a href="#">Q07817</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user