

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

ACTH (Adrenocorticotrophic Hormone) (C-Terminus) (CLIP/1449), CF647 conjugate, 0.1mg/mL, IgG1, Clone: [CLIP/1449], Mouse, Monoclonal BOT-BNC471449-100

| | |
|----------------------------|---|
| Article Name | ACTH (Adrenocorticotrophic Hormone) (C-Terminus) (CLIP/1449), CF647 conjugate, 0.1mg/mL, IgG1, Clone: [CLIP/1449], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC471449-100 |
| Supplier Catalog Number | BNC471449-100 |
| Alternative Catalog Number | BOT-BNC471449-100-100UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Synthetic peptide corresponding to aa25-39 of human ACTH (NGAEDESAEAFPLEF) |
| Conjugation | CF647 |
| Product Description | ACTH (same as Corticotropin) is a 39 amino acid active peptide produced by the anterior pituitary. This MAb is specific to CLIP (aa25-39 of ACTH), does not react with Synacthen (aa1-24 of ACTH). POMC (pro-opiomelanocortin or corticotropin-lipotropin)... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |

| | |
|-------------------|---|
| Clone Designation | [CLIP/1449] |
| Molecular Weight | ACTH is ~5 kDa |
| Isotype | IgG1 |
| UniProt | P01189 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL ELISA For coating, order Ab without BSA Optimal dilution for a specific application should be determined by user |