

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Bcl-6 (Follicular Lymphoma Marker) (BCL6/1527), CF647 conjugate, 0.1mg/mL, Clone: [BCL6/1527], Mouse, Monoclonal BOT-BNC471527-500**

Article Name	Bcl-6 (Follicular Lymphoma Marker) (BCL6/1527), CF647 conjugate, 0.1mg/mL, Clone: [BCL6/1527], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC471527-500
Supplier Catalog Number	BNC471527-500
Alternative Catalog Number	BOT-BNC471527-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant human bcl-6 protein fragment (aa256-389) (exact sequence is proprietary)
Conjugation	CF647
Product Description	Bcl-6 is a transcriptional repressor mainly required for germinal center (GC) formation and antibody affinity maturation which has different mechanisms of action specific to the lineage and biological functions. It forms complexes with different core...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[BCL6/1527]

Molecular Weight	95 kDa
UniProt	<a href="#">P41182</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user