

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Cytokeratin, pan (Epithelial Marker) (KRTH/1576R + KRTL/1577R), CF647 conjugate, 0.1mg/mL, Clone: [KRTH/1576R KRTL/1577R], Rabbit, Monoclonal BOT-BNC471600-100**

|                            |   |
|----------------------------|---|
| Article Name               | Cytokeratin, pan (Epithelial Marker) (KRTH/1576R + KRTL/1577R), CF647 conjugate, 0.1mg/mL, Clone: [KRTH/1576R KRTL/1577R], Rabbit, Monoclonal   |
| Biozol Catalog Number      | BOT-BNC471600-100   |
| Supplier Catalog Number    | BNC471600-100   |
| Alternative Catalog Number | BOT-BNC471600-100-100UL   |
| Manufacturer               | Biotium   |
| Host                       | Rabbit  |
| Category                   | Antikörper  |
| Application                | IHC   |
| Species Reactivity         | Human   |
| Immunogen                  | Recombinant full-length human KRT76 and KRT77 proteins  |
| Conjugation                | CF647   |
| Product Description        | Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI <5.7) and basic (pI >6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, which 67 kDa (CK1), 6... |
| Clonality                  | Monoclonal  |
| Concentration              | 0.1 mg/mL   |

|                   |  |
|-------------------|--|
| Clone Designation | [KRTH/1576R KRTL/1577R]  |
| Molecular Weight  | 40-67 kDa  |
| UniProt           | <a href="#">Q7Z794</a>   |
| Buffer            | PBS, 0.1% BSA, 0.05% azide   |
| Source            | Animal   |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution for a specific application should be determined by user |