

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **CD31 / PECAM-1 (Endothelial Cell Marker)(1A10), CF568 conjugate, 0.1mg/mL, Clone: [1A10], Mouse, Monoclonal BOT-BNC680846-100**

Article Name	CD31 / PECAM-1 (Endothelial Cell Marker)(1A10), CF568 conjugate, 0.1mg/mL, Clone: [1A10], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC680846-100
Supplier Catalog Number	BNC680846-100
Alternative Catalog Number	BOT-BNC680846-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Species Reactivity	Human
Immunogen	Prokaryotic recombinant fusion protein corresponding to a portion of the extracellular domain downstream of the signal sequence of the CD31 (PECAM-1)
Conjugation	CF568
Product Description	CD31 (PECAM-1) is a transmembrane glycoprotein member of the immunoglobulin supergene family of adhesion molecules. CD31 is expressed by stem cells of the hematopoietic system and is primarily used to identify and concentrate these cells for experime...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[1A10]

Molecular Weight	~100 kDa (endothelium), ~130 kDa (platelets)
UniProt	<a href="#">P16284</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry