

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Mucin 5AC (MUC5AC)(MUC5AC/917), CF568 conjugate, 0.1mg/mL, Clone: [MUC5AC/917], Mouse, Monoclonal BOT-BNC680917-100

| | |
|----------------------------|---|
| Article Name | Mucin 5AC (MUC5AC)(MUC5AC/917), CF568 conjugate, 0.1mg/mL, Clone: [MUC5AC/917], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC680917-100 |
| Supplier Catalog Number | BNC680917-100 |
| Alternative Catalog Number | BOT-BNC680917-100-100UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant human MUC5AC protein |
| Conjugation | CF568 |
| Product Description | Mucin 5AC glycoprotein (MUC5AC) is a 641 kDa glycoprotein belonging to the superfamily of mucins. Mucins are high molecular weight glycoproteins produced by epithelial cells and can be divided into two families, secretory mucins and membrane bound mu... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [MUC5AC/917] |

| | |
|-------------------|---|
| Molecular Weight | 641 kDa |
| UniProt | P98088 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology formalin-fixed 1-2 ug/mL Does not react with rat, others not known Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |