

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **UACA / Nucling(AE-5), CF568 conjugate, 0.1mg/mL, Clone: [AE-5], Mouse, Monoclonal BOT-BNC680956-500**

|                            |   |
|----------------------------|---|
| Article Name               | UACA / Nucling(AE-5), CF568 conjugate, 0.1mg/mL, Clone: [AE-5], Mouse, Monoclonal   |
| Biozol Catalog Number      | BOT-BNC680956-500   |
| Supplier Catalog Number    | BNC680956-500   |
| Alternative Catalog Number | BOT-BNC680956-500-500UL   |
| Manufacturer               | Biotium   |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | WB  |
| Species Reactivity         | Human, Mouse  |
| Immunogen                  | Nuclei of myeloid leukemia biopsy cells   |
| Conjugation                | CF568   |
| Product Description        | UACA (Uveal Autoantigen with Coiled-coil domains and Ankyrin repeats) is a 1,416 amino acid nuclear membrane protein. It was originally identified as an autoantigen in patients with panuveitis, a characteristic of Vogt-Koyanagi-Harada disease, and in... |
| Clonality                  | Monoclonal  |
| Concentration              | 0.1 mg/mL   |
| Clone Designation          | [AE-5]  |

|                   |   |
|-------------------|---|
| Molecular Weight  | 160 kDa   |
| UniProt           | <a href="#">Q9BZF9</a>  |
| Buffer            | PBS, 0.1% BSA, 0.05% azide  |
| Source            | Animal  |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 0.5-1.0 ug/mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |