

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD63(LAMP3/968), CF568 conjugate, 0.1mg/mL, Clone: [LAMP3/968], Mouse, Monoclonal BOT-BNC680968-500

| | |
|----------------------------|---|
| Article Name | CD63(LAMP3/968), CF568 conjugate, 0.1mg/mL, Clone: [LAMP3/968], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC680968-500 |
| Supplier Catalog Number | BNC680968-500 |
| Alternative Catalog Number | BOT-BNC680968-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | Exosome Staining, FC, IF, IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant human full-length CD63 protein |
| Conjugation | CF568 |
| Product Description | This MAb recognizes protein of 26 kDa-60 kDa, which is identified as CD63. Its epitope is different from that of MAb LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells an... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [LAMP3/968] |

| | |
|-------------------|--|
| Molecular Weight | 26 kDa (core protein), 30-60 kDa (glycosylated) |
| UniProt | P08962 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 1-2 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |